

# Pediatric Physicians, PC

## IBUPROFEN

- Choose Advil, Motrin, or generic ibuprofen
- Do NOT use ibuprofen products in babies less than 6 months
- Ibuprofen may be given every 6 hours

WEIGHT (pounds)	ORAL DROPS 50mg/1.25ml	SUSPENSION 100mg/5ml	CHEWABLE TABLETS 50mg	CHEWABLE TABLETS 100mg	JUNIOR CAPLETS 100mg	CAPSULES 200mg
12-17	1.25 ml	½ tsp				
18-23	1.875 ml	¾ tsp	1 ½ tab			
24-35	2.5 ml	1 tsp	2 tab	1 tab		
36-47		1 ½ tsp	3 tab	1 ½ tab		
48-59		2 tsp	4 tab	2 tab	2 cap	1 cap
60-71		2 ½ tsp	5 tab	2 ½ tab	2 ½ cap	1 cap
72-95		3 tsp	6 tab	3 tab	3 cap	2 cap
> 95		4 tsp	8 tab	4 tab	4 cap	2 cap

## ACETAMINOPHEN

- Choose Tylenol or generic acetaminophen
- Acetaminophen may be given every 4-6 hours, not to exceed 5 doses in 24 hours
- Infants' acetaminophen concentrated drops, 80 mg/.8 ml with built-in dropper, were discontinued in 2011. Infants can use infants' suspension.
- Always use the dosing device that came with the package, and follow label instructions.

WEIGHT POUNDS	INFANTS' CONCENTRATED DROPS (has built-in dropper) 80 mg/0.8 mL	CHILDREN'S or INFANTS' SUSPENSION 160 mg/5 mL	CHILDREN'S CHEW TABS 80 mg each	JR STRENGTH CAPS 160 mg each	ADULT CAPS 325 mg each
6-11	.4 mL	1.25 mL			
12-17	.8 mL	2.5 mL			
18-23	1.2 mL	3.75 mL			
24-35	1.6 mL	5 mL	2 tab		
36-47		7.5 mL	3 tab		
48-59		10 mL	4 tab	2 tab	1 tab
60-71		12.5 mL	5 tab	2 ½ tab	1 tab
72-up		15 mL	6 tab	3 tab	2 tab

For more information, visit [www.PediatricPhysiciansPC.com](http://www.PediatricPhysiciansPC.com)