

Pediatric Physicians, PC

IBUPROFEN

- Choose Advil, Motrin, or generic ibuprofen
- Do NOT use ibuprofen products in babies less than 6 months
- Ibuprofen may be given every 6 hours

| WEIGHT (pounds) | ORAL DROPS 50mg/1.25ml | SUSPENSION 100mg/5ml | CHEWABLE TABLETS 50mg | CHEWABLE TABLETS 100mg | JUNIOR CAPLETS 100mg | CAPSULES 200mg |
|-----------------|---------------------------|-------------------------|--------------------------|---------------------------|-------------------------|-------------------|
| 12-17 | 1.25 ml | ½ tsp | | | | |
| 18-23 | 1.875 ml | ¾ tsp | 1 ½ tab | | | |
| 24-35 | 2.5 ml | 1 tsp | 2 tab | 1 tab | | |
| 36-47 | | 1 ½ tsp | 3 tab | 1 ½ tab | | |
| 48-59 | | 2 tsp | 4 tab | 2 tab | 2 cap | 1 cap |
| 60-71 | | 2 ½ tsp | 5 tab | 2 ½ tab | 2 ½ cap | 1 cap |
| 72-95 | | 3 tsp | 6 tab | 3 tab | 3 cap | 2 cap |
| > 95 | | 4 tsp | 8 tab | 4 tab | 4 cap | 2 cap |

ACETAMINOPHEN

- Choose Tylenol or generic acetaminophen
- Acetaminophen may be given every 4-6 hours, not to exceed 5 doses in 24 hours
- Always use the dosing device that came with the package, and follow label instructions.

| WEIGHT POUNDS | CHILDREN'S or INFANTS' SUSPENSION 160 mg/5 mL | CHILDREN'S CHEW TABS 80 mg each | JR STRENGTH CAPS 160 mg each | ADULT CAPS 325 mg each |
|---------------|--|------------------------------------|---------------------------------|---------------------------|
| 6-11 | 1.25 mL | | | |
| 12-17 | 2.5 mL | | | |
| 18-23 | 3.75 mL | | | |
| 24-35 | 5 mL | 2 tab | | |
| 36-47 | 7.5 mL | 3 tab | | |
| 48-59 | 10 mL | 4 tab | 2 tab | 1 tab |
| 60-71 | 12.5 mL | 5 tab | 2 ½ tab | 1 tab |
| 72-up | 15 mL | 6 tab | 3 tab | 2 tab |